

# produce feed, fuel and fertilizer from green biomass – An organic biorefinery concept

M. Santamaría, B. Molinuevo-Salces<sup>1</sup>, M. Lübeck and H. Uellendahl<sup>\*</sup>.

Section for Sustainable Biotechnology, Aalborg University Copenhagen, A.C. Meyers Vænge 15, 2450 København, Denmark.

<sup>\*</sup>Corresponding author. Tel. +45 9940 2585, E-mail address: [hu@bio.aau.dk](mailto:hu@bio.aau.dk)

## ORGANOFINERY CONCEPT – SEPARATION EFFICIENCIES

- ✓ **Press cake and brown juice** after protein extraction accounted for more than 90% of fresh biomass with 61-81% of VS found in the press cake and 7-11% of VS in the brown juice – energy potential for biogas production.
- ✓ Recovery of crude proteins in the **protein concentrate** was more than 15%.





Figure 1. Separation of total mass, VS and crude protein into the output fractions.


## THE PROTEIN CONCENTRATE – ORGANIC POULTRY FEED

- ✓ Crude protein content in the concentrate is around 40% (dry matter basis).
- ✓ Amino acid profile of **protein concentrate** comparable with current organic poultry feed.




## SPECIFIC METHANE YIELDS

- ✓ The specific methane yield of the **press cake** from red clover and clover grass was 70% and 90% of the fresh biomass (331 and 344 ml-CH<sub>4</sub>/g-VS), respectively).
- ✓ The **press cake** presented enough nutrients to ensure a successful anaerobic digestion.

Figure 2. Specific methane yield (mL-CH<sub>4</sub>/g-VS) during 55 days of AD.

- ✓ High specific methane yield of **brown juice** due to high content of easily degradable organic matter. Low pH (4-5), may need pH control.
- ✓ Some substrate inhibition at high substrate loading.

Figure 3. Specific methane yield (mL-CH<sub>4</sub>/g-VS) of brown juice after 15 and 55 days of anaerobic digestion for 2 different substrate loadings (S/l ratio of 1 and 0.5).

## BIOGAS POTENTIAL RECOVERY

- ✓ Up to 85% of the methane potential of the fresh crops is still preserved in the residual fractions after protein refining (Fig. 4).
  - 41-69% of the potential methane of the fresh crop is recovered in the **press cake**.
  - 10-16% of the potential methane of the fresh crop is recovered in the **brown juice**.
- ✓ The digestate will be used as organic fertilizer.




Figure 4. Methane yield of the fresh biomass (m³/t) and of the residual biomass fractions after the biorefinery process (m³/t of fresh crop).

<sup>1</sup>Present address: IMDEA Energy, Avda. Ramón de la Sagra 3, 28935 Móstoles, Madrid (Spain).

**Acknowledgements:** The Organofinery project is supported from "Green Development and Demonstration Program (GUDP) under the Danish Ministry of Food" and it is part of the Organic RDD-2 program, coordinated by ICROFS.